

Evaluation of nutritional status of patients with gastrointestinal system tumours receiving chemo-radiotherapy

Nutrition in patients receiving chemo-radiotherapy

Nazan Son¹, Ezgi Atıcı²

¹ Department of Nutrition and Dietetics, Faculty of Health Sciences, Anadolu University, Eskisehir

² Department of Nutrition and Dietetics, Ceylan Hospital, Bursa, Turkey

Abstract

Aim: This study was planned to examine the nutritional problems of patients diagnosed with gastrointestinal tract (GIS) cancer and to evaluate the effects of professional nutritional support on the outcome of their treatment.

Material and Methods: This study was carried out with 24 (33.3% females, 66.7% males) volunteer participants aged 18 and above, who were diagnosed with GIS cancer and planned to have concurrent chemotherapy and radiotherapy treatment (chemoradiotherapy=CRT). The participants were asked, questions on socio-demographic characteristics and nutritional habits, anthropometric measurements were taken, biochemical data were recorded, Subjective Global Assessment (SGA) and Mini Nutritional Assessment (MNA) scales were applied, and nutritional training deemed necessary were given to them. The participants were re-evaluated after three months and the data were compared with the initial findings.

Results: There was no significant difference between the pre-and post-treatment Body Mass Index (BMI). A significant difference was found between the SGD and MNA values before and after CRT ($p=0.000$).

Discussion: In cancer patients, alongtogether with the medical treatment plan, the patient's nutritional evaluation and providing giving the necessary Medical Nutrition Therapy (MNT) enable the individual to act consciously. Increasing the level of nutritional awareness makes it easier for an individual to fight the disease while increasing the effectiveness of treatment. In conclusion, it should be noted that a dietitian should be available in institutions providing care to cancer patients should have a dietitian.

Keywords

Gastrointestinal Cancer, Nutrition, Malnutrition, Subjective Global Assessment, Mini Nutritional Assessment.

DOI: 10.4328/ACAM.22006 Received: 2023-10-05 Accepted: 2023-11-06 Published Online: 2023-11-14 Printed: 2024-01-01 Ann Clin Anal Med 2024;15(1):47-51

Corresponding Author: Nazan Son, Department of Nutrition and Dietetics, Faculty of Health Sciences, Anadolu University, Eskisehir, Turkey.

E-mail: nazanson@gmail.com P: +90 505 448 14 41

Corresponding Author ORCID ID: <https://orcid.org/0000-0003-3614-3604>

This study was approved by the Ethics Committee of Okan University (Date: 2021-01-27, No: 132)

Introduction

Cancer is among the most important health problems all over the world. It is known to be the second leading among the causes of death in the world [1-3]. The causes of cancer can be grouped under two headings as environmental and genetic causes. For example; Smoking, consumed food and beverages, obesity, hormones, viruses, physical and chemical agents are some of the environmental factors. [4]. According to data from the International Agency for Research on Cancer in 2018, 18.1 million people have been diagnosed with cancer worldwide, and it is estimated that 24 million people will have cancer in 2035 [3]. According to Globocan 2020 data, GIS cancers are the leading cause of cancer-related death. In 2020, 5 million new GIS cancer diagnoses were causeding approximately 3.5 million deaths [4]. While the population of Turkey was 84.339,067 in 2020, a total of 233,834 new cancer cases were detected in the same year. Of this number, 126,335 resulted in death [5].

There are five main treatment options in cancer treatment: chemotherapy, radiotherapy, surgical treatment, immunotherapy, and hormone therapy. In recent years, the application of concomitant CRT has become an important practice in cancer treatment. This practice has become a standard treatment for advanced cancer types [6]. Many cancer treatments affect patients' appetite, food intake, and the use/absorption of nutrients in the body. Especially in patients who have undergone gastrointestinal surgery, malabsorption of nutrients develops since swallowing and gastrointestinal motility are affected [7]. The purpose of MNT in cancer patients is to prevent and treat nutritional deficiencies, reduce the side effects of treatment, prevent weight loss, support the effectiveness of cancer treatment, reduce the rate of complications and mortality, and improve quality of life [8]. Personalized nutrition training has been reported to increase the patient's food consumption and quality of life, and reduce the complications related to the disease in cancer patients who cannot have enough nutrients [4,7]. Therefore, this study was conducted to evaluate the effectiveness and necessity of MNT given to patients who have been diagnosed with GIS cancer before starting CRT treatment.

Material and Methods

The population of this prospective cohort study consisted of patients who were newly diagnosed with GIS cancer in Bursa Ali Osman Sönmez Oncology Hospital between 01 March 2021 and 01 June 2021. The study was carried out with a total of 24 patients, 8 females (33.3%) and 16 males (66.7%) aged 18 years and older, who were diagnosed with GIS cancer and were scheduled for CRT treatment. Before the treatment, all patients were given nutritional training and nutritional follow-up of the patients was carried out. Patients over 85 years of age and those with the presence of metastases were not included in the study.

The same Dietitian has given nutrition training to the participants individually before the treatment and their nutrition follow-ups were carried out.

Socio-demographic Data Collection Form included questions about gender, age, weight, height, education level, occupation, monthly income, presence of additional disease and physical activity levels of the patients, and their BMI values were

calculated and evaluated with the data obtained from this form. Eating Habits Questionnaire Form included questions about the number of main and snack meals consumed by the patients, whether they skipped meals, what type of food they consumed between main meals, and the frequency of eating outside the home in order to evaluate the nutritional status of individuals. Furthermore, the three-day food consumption of the patients was recorded with the form created and evaluated by the same dietitian.

Anthropometric measurements of the participants were taken as per the standards. The body weight (kg), height (cm), triceps skinfold thickness (TSF), mid-upper arm circumference (MUAC), and Midarm Muscle Area (MMA) measurements of the participants were taken and recorded by the researcher.

Collection of biochemical parameters: Total protein, Albumin, Hemoglobin, total cholesterol, HDL cholesterol (mg/dL), LDL-cholesterol (mg/dL), Triglycerides, C-reactive protein (CRP), CEA, CA19-9, and white blood cell (WBC), SGOT(BOTTOM), SGPT(AST), fasting blood glucose, urea, creatinine, calcium, iron and vitamin B12 values of the patients have been obtained from the patient files and recorded.

Scales applied to participants:

The MNA was developed by Vellas et al. [9]. The Turkish validity and reliability study of the scale was carried out by Sarıkaya in 2013 [10].

The SGD was developed by Detsky et al. to assess malnutrition status. It consists of two parts: dietary history and physical examination. In the scoring of the scale, SGD-A indicates normal nutritional status, SGD-B suspected malnutrition and SGD-C severe malnutrition [11].

Statistical analysis of data:

Analysis of the data was performed with the SPSS 25.0.0 program, while numerical variables were measured with mean \pm standard deviation and lower-higher, categorical variables were shown as numbers and percentages. "Wilcoxon signed-rank test" was used for quantitative initial measurement and final measurement comparisons. The level of statistical significance was determined as $p < 0.05$.

Ethics Committee Approval

Ethics Committee approval for the study was obtained on 27.01.2021 from the Ethics Committee of Okan University.

Results

Of the 24 study participants in the study, 33% were female. The mean age was 60.13 ± 11.72 years. The educational status of the participants was as follows: 45.8% were primary school graduates, while 20.8% were university graduates. The monthly incomes of 12.5% of the participants were 2500 TL or less, 45.8% had a monthly income of were between 2501-5000 TL, 33.3%: were between 5001-10000 TL, and 8.3%: were 10001 TL and above. Among the participants, 45.8% of them did not have any chronic disease. When the daily physical activity levels were examined, 54.2% of the participants were found to perform physical activity. 16.7% of the participants actively smoked and 20.8% of social drinkers consumed alcohol. A total of 70.8% of the participants had an immediate family member who was diagnosed with cancer. The information about the nutritional habits of the participants is as follows.

While 50% of the participants have had two main meals, 50% have had three main meals a day. Of these, A total of 37.5% of them hadve one snack a day, 25% had two and 8.3% had three snacks. While 25% of the participants preferred candy, chocolate and wafers as snacks, 16.7% of them consumed bagels, biscuits, and cookies and 12.5% of them consumed carbonated drinks and soda water. Ready-made food is was consumed by 41.7% of the participants.

The findings regarding the comparison of the anthropometric characteristics of the participants before and after CRT are presented given in Table 2. There was no statistically significant difference between the body weight and mean BMI values of the participants (p>0.05).

The only statistically significant difference was found between the midarm muscle circumference values of the participants before CRT and after CRT (p=0.016).

There was no statistically significant difference in the biochemical parameters of the participants before and after

Table 1. Distribution of participants in SGD and MNA groups before and after CRT

	Before CRT n (%)				P	After CRT n (%)				P
	SGD -A	SGD -B	SGD -C	Total		SGD -A	SGD -B	SGD -C	Total	
Malnutrition	0 (0.0)	1 (4.2)	2 (8.2)	3 (12.5)		0 (0.0)	0 (0.0)	2 (8.3)	2 (8.3)	
Risk of Malnutrition	2 (8.3)	6 (25.0)	3 (12.5)	11 (45.8)		2 (8.3)	7 (29.2)	0 (0.0)	9 (37.5)	
Normal Nutrition	10 (41.7)	0 (0.0)	0 (0.0)	10 (41.7)		11 (45.8)	2 (8.3)	0 (0.0)	13 (54.2)	
Total	12 (50.0)	7 (29.2)	5 (29.2)	24 (100)		13 (54.2)	9 (37.5)	2 (8.3)	24 (100)	

Table 2. Comparison of the anthropometric characteristics of the participants before and after CRT

n=24	Before CRT x±Sd min-max	After CRT x±Sd min-max	P
Weight (kg)	67.33±11.23 50-96	66.58±9.04 54-86	0.666
Height(cm)	162.86±35.14 1.56-183	162.77±35.03 1.56-183	1.000
BMI (kg/m ²)	23.66±4.56 15.8-37.5	23.88±3.48 18.5-33.6	0.455
Triceps Skinfold Thickness (TST) (mm)	16.25±9.13 5-36	16.00±7.39 5-30	0.972
Mid-Upper Arm Circumference (MUAC) (cm)	29.75±4.01 23-37	29.33±3.90 22-35	0.545
Midarm Muscle Circumference (MMC) (cm)	25.71±7.36 18-56	29.58±13.94 18-72	0.016
Midarm Muscle Area (MMA) (cm ²)	48.91±14.44 26.60-74	49.84±14.27 29-74	0.274

Table 3. Correlation of participants' daily calorie intakes, BMI, and MNA values before and after CRT

		Before CRT		After CRT	
		BMI	MNA	BMI	MNA
Daily Calorie	R	0.027	0.147	0.015	0.254
	P	0.899	0.493	0.943	0.231
BMI	R	1.000	0.386	1.000	0.349
	P		0.063		0.095
MNA	R	0.386	1.000	0.349	1.000
	P	0.063		0.095	

CRT, except for the ferritin value (p=0.026).

As per the SGD, the results of the participants before CRT were found to be 50% SGD-A, 29.2% SGD-B, 20.8% SGD-C, and after CRT, 54.2% of them were found to be SGD-A, % 37.5% SGD-B, and 8.3% to SGD-C. According to the MNA, before CRT, 12.5% of the participants were found to have malnutrition, 45.8% were at risk of malnutrition, 41.7% were in the normal nutrition group, while 8.3% had malnutrition, 37.5% were at risk of malnutrition and 54.2% of them had normal nutrition after CRT. A significant difference was found between the SGD and MNA values of the participants before and after CRT (p=0.000). (Table 1).

Correlation of participants' daily calorie intakes, BMI, and MNA values before and after CRT, Nno significant relationship was found between the daily calorie intake, BMI and MNA levels of the participants before and after CRT (p>0.05) (Table 3).

Discussion

Malnutrition, poor nutrition and muscle loss are frequently observed in cancer patients. This condition negatively affects the clinical results of patients and leads to an increase in the length of their stay in the hospital [12]. This study was planned to evaluate how nutritional status before and after CRT in individuals with GIS cancer and professional support affect the treatment.

Some studies have reported that gender-specific cancer It has been reported in some studies that the incidence of cancer in terms of gender in Turkey is higher in males than in females [13]. In some studies conducted in different countries of the world, the incidence of cancer has also been reported also to be higher in males compared to females [14-15]. The fact that the majority of the participants in our study were male supports this literature finding. There are studies reporting that low socioeconomic [14,16] and low educational status [16] trigger the incidence of cancer. Our results have been in line with this information.

It is noted in the literature that patients with a family history of gastrointestinal cancer tend to be diagnosed with cancer at a younger age than those without [15]. Among our study the participants of our study, 70.8% of them had an immediate family member who had been diagnosed with a type of cancer. Half of the participants in our study stated that they skipped a main meal daily. Similar to our study results, two different studies reported that almost half of the participants skipped one of the main meals [17]. But, in our study, the daily calorie consumption and macronutrient distribution of the participants before and after CRT were similar, and no statistically significant difference was found (p>0.05).

Malnutrition is closely associated with a worse prognosis and poor quality of life [18]. In the ir study by Demirel et al. conducted with patients with head and neck cancer in our country, Demirel et al. they found a significant relationship between SGD and MNA (p<0.001) [19].

In our study, the mean MNA score of the participants before CRT was 21.38±4.30, while it was 23.42±3.67 after CRT, and a significant difference was found between the scores (p=0.008). This changing situation in favour of the participants shows that individuals consume food consciously to meet their daily

energy needs, despite the side effects that may develop due to CRT. This also shows us that patients care about nutritional advice about coping with the disease, and they act diligently and consciously on their own. It also indicates that they have been in compliance with the treatment. The results of our study emphasize the importance of MNT.

In the study conducted by Hong et al. with the participants who were treated for cancer, they reported finding lower TST values in male participants, and lower MUAC, MMC, and MMA values in both male and female participants compared to the baseline values [20]. In our study, while the MUAC value of the participants did not change, the MMC and MMA values increased (Table 2). This was believed to be due to the fact that the participants were able to continue their daily activities as a result of their compliance with MNT and feeling well during the treatment process.

Biochemical Characteristics of Patients Before and After CRT:

Many biochemical parameters need to be evaluated before a malnutrition diagnosis can be made. It can play an important role in the diagnosis of the disease at an early stage [21].

In the study of by Kabata et al., the albumin value of the participants was 3.8, total protein value was 7 (4.5-7.9) [22]; Lee et al. was found that total protein was 5.8 (4.5-7.9) [23]. In our study, total protein it was found to be total protein 63.95 ± 28.19 (64-83) before CRT, while it was 64.92 ± 10.15 after CRT.

It was proven in a meta-analysis study that a low haemoglobin value causes a poor prognosis [24]. In our study, the haemoglobin value of the participants was found to be higher than the value before CRT. The CRP level of the participants was found to be high in our study. Inflammation in some tissues with the development of cancer or an increase in CRP level may be observed as the body's immune response [24].

In another study conducted with a total of 4707 patients and Huang et al. Were associated with inflammatory markers ferritin and CRP [25]. Apart from CRT, the level of iron in the blood is also affected by liver diseases and inflammation [1]. There was a statistically significant difference only in the ferritin value among the biochemical parameters of the participants before and after CRT in our study ($p=0.026$). This difference was in favour of the patients and in the direction of increase. No significant difference was observed in other data.

Conclusion:

our study results once again revealed the importance of MNT in cancer patients. Having a Nutritionist in the units providing oncology healthcare services and giving the necessary MNT to the patients would not only enable faster improvements in the health parameters of the patients but would also increase their quality of life. It would also contribute to the economy of the countries by reducing treatment costs.

Scientific Responsibility Statement

The authors declare that they are responsible for the article's scientific content including study design, data collection, analysis and interpretation, writing, some of the main line, or all of the preparation and scientific review of the contents and approval of the final version of the article.

Animal and Human Rights Statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with

the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Funding: None

Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Conti CB, Agnesi S, Scaravaglio M, Masseria P, Dinelli ME, Oldani M, et al. Early Gastric Cancer: Update on Prevention, Diagnosis and Treatment. *Int J Environ Res Public Health*. 2023;20(3):2149.
2. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018, GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394-424.
3. Weeden CE, Hill W, Lim EL, Grönroos E, Swanton C. Impact of risk factors on early cancer evolution. *Cell*. 2023;186(8):1541-63.
4. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021; 71(3):209-49.
5. Topakta B, Karakus B, Yaksi N. Clinical and epidemiological characteristic of cancer cases reported between 2013-2017 in Amasya. *Turk J Public Health*. 2023;21(1):113-26.
6. Wang Y, Shen L, Wan J, Zhang H, Wu R, Wang J, et al. Neoadjuvant chemoradiotherapy combined with immunotherapy for locally advanced rectal cancer: A new era for anal preservation. *Front Immunol*. 2022;13:1067036.
7. Li B, Shao H, Gao L, Li H, Sheng H, Zhu L. Nano-drug co-delivery system of natural active ingredients and chemotherapy drugs for cancer treatment: a review. *Drug Deliv*. 2022;29(1):2130-61.
8. Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr*. 2017;36(1):11-48.
9. Vellas B, Guigoz Y, Garry PJ, Nourhashemi F, Bennahum D, Lauque S, et al. The Mini Nutritional Assessment (MNA) and its Use in Grading the Nutritional State of Elderly Patients. *Nutrition*. 1999;15(2):116-22.
10. Sarıkaya D, Halil M, Kuyumcu ME, Kilic MK, Yesil Y, Kara O, et al. Mini nutritional assessment test long and short form are valid screening tools in Turkish older adults. *Archives of Gerontology and Geriatrics*. 2015;61(1):56-60.
11. Detsky AS, McLaughlin JR, Baker JP, Johnston N, Whittaker S, Mendelson RA, et al. What is subjective global assessment of nutritional status? *J Parenter Enteral Nutr*. 1987;11(1):8-13.
12. Castillo-Martínez L, Castro-Eguiluz D, Copca-Mendoza ET, Pérez-Camargo DA, Reyes-Torres CA, Ávila EA, et al. Nutritional Assessment Tools for the Identification of Malnutrition and Nutritional Risk Associated with Cancer Treatment. *Rev Invest Clin*. 2018;70(3):121-5.
13. Huang J, Ngai CH, Deng Y, Pun CN, Lok V, Zhang L, et al. Incidence and mortality of thyroid cancer in 50 countries: a joinpoint regression analysis of global trends. *Endocrine*. 2023;80:355-65.
14. Pang Y, Kartsonaki C, Guo Y, Chen Y, Yang L, Bian Z, et al. Socioeconomic Status in Relation to Risks of Major Gastrointestinal Cancers in Chinese Adults: A Prospective Study of 0.5 Million People. *Cancer Epidemiol Biomarkers Prev*. 2020;29(4):823-31.
15. Chung JW, Park JJ, Lim YJ, Lee J, Kim SM, Han JH, et al. Gastrointestinal cancer risk in patients with a family history of gastrointestinal cancer. *Korean J Gastroenterol*. 2018;71(6):338-48.
16. Kawakatsu Y, Koyanagi YN, Oze I, Kasugai Y, Morioka H, Yamaguchi R, et al. Association between Socioeconomic Status and Digestive Tract Cancers: A Case-Control Study. *Cancers*. 2020;12(11):3258.
17. Gunes-Bayir A, Kiziltan HS, Sentürk N, Mayadagli A, Gumus M. A pilot study of self-reported physical activity and eating habits in Turkish Cancer patients under chemotherapy. *Nutrition and Cancer*. 2015;67(6):906-11.
18. Maia FCP, Silva TA, Generoso SV, Correia MITD. Malnutrition is associated with poor health-related quality of life in surgical patients with gastrointestinal cancer. *Nutrition*. 2020;75-76:110769.
19. Demirel B, Atasoy BM. Comparison of Three Nutritional Screening Tools to Predict Malnutrition Risk and Detect Distinctions Between Tools in Cancer Patients Receiving Radiochemotherapy. *Nutr Cancer*. 2018;70(6):867-73.
20. Hong JS, Wu LH, Su L, Zhang HR, Lv WL, Zhang WJ, et al. Effect of chemoradiotherapy on nutrition status of patients with nasopharyngeal cancer. *Nutr Cancer*. 2016;68(1):63-9.
21. Jabłońska B, Mrowiec S. Nutritional Status and Its Detection in Patients with Inflammatory Bowel Diseases. *Nutrients*. 2023;15(8):1991.
22. Kabata P, Jastrzębski T, Kąkol M, Król K, Bobowicz M, Kosowska A, et al. Preoperative nutritional support in cancer patients with no clinical signs of malnutrition-prospective randomized controlled trial. *Supportive Care in Cancer*. 2015;23(2):365-70.
23. Lee SY, Lee YJ, Yang JH, Kim CM, Choi WS. The association between phase angle of bioelectrical impedance analysis and survival time in advanced cancer patients: preliminary study. *Korean J Fam Med*. 2014;35(5):251-6.
24. Huang XZ, Yang YC, Chen Y, Wu CC, Lin RF, Wang ZN, et al. Preoperative Anemia or Low Hemoglobin Predicts Poor Prognosis in Gastric Cancer Patients: A Meta-Analysis. *Dis Markers*. 2019;7606128.
25. Oztas D, Erdogan S, Benzil A, Nadar O, Erel O. A New Inflammatory Marker: Vitamin B12. *J Immunol Clin Microbiol*. 2019;4(1):1-12.

How to Cite This Article:

Nazan Son, Ezgi Atıcı. Evaluation of nutritional status of patients with gastrointestinal system tumours receiving chemo-radiotherapy. *Ann Clin Anal Med* 2024;15(1):47-51

This study was approved by the Ethics Committee of Okan University (Date: 2021-01-27, No: 132)